

RECENT INSIGHTS INTO THE DOUBLE ROLE OF HYDROGEN PEROXIDE IN PLANTS

EDITED BY: Naser A. Anjum, Sarvajeet Singh Gill, Francisco J. Corpas,
Cristina Ortega-Villasante, Luis E. Hernandez, Narendra Tuteja,
Adriano Sofo, Mirza Hasanuzzaman and Masayuki Fujita

PUBLISHED IN: *Frontiers in Plant Science*





frontiers

Frontiers eBook Copyright Statement

The copyright in the text of individual articles in this eBook is the property of their respective authors or their respective institutions or funders. The copyright in graphics and images within each article may be subject to copyright of other parties. In both cases this is subject to a license granted to Frontiers.

The compilation of articles constituting this eBook is the property of Frontiers.

Each article within this eBook, and the eBook itself, are published under the most recent version of the Creative Commons CC-BY licence.

The version current at the date of publication of this eBook is CC-BY 4.0. If the CC-BY licence is updated, the licence granted by Frontiers is automatically updated to the new version.

When exercising any right under the CC-BY licence, Frontiers must be attributed as the original publisher of the article or eBook, as applicable.

Authors have the responsibility of ensuring that any graphics or other materials which are the property of others may be included in the CC-BY licence, but this should be checked before relying on the CC-BY licence to reproduce those materials. Any copyright notices relating to those materials must be complied with.

Copyright and source acknowledgement notices may not be removed and must be displayed in any copy, derivative work or partial copy which includes the elements in question.

All copyright, and all rights therein, are protected by national and international copyright laws. The above represents a summary only. For further information please read Frontiers' Conditions for Website Use and Copyright Statement, and the applicable CC-BY licence.

ISSN 1664-8714

ISBN 978-2-88974-524-1

DOI 10.3389/978-2-88974-524-1

About Frontiers

Frontiers is more than just an open-access publisher of scholarly articles: it is a pioneering approach to the world of academia, radically improving the way scholarly research is managed. The grand vision of Frontiers is a world where all people have an equal opportunity to seek, share and generate knowledge. Frontiers provides immediate and permanent online open access to all its publications, but this alone is not enough to realize our grand goals.

Frontiers Journal Series

The Frontiers Journal Series is a multi-tier and interdisciplinary set of open-access, online journals, promising a paradigm shift from the current review, selection and dissemination processes in academic publishing. All Frontiers journals are driven by researchers for researchers; therefore, they constitute a service to the scholarly community. At the same time, the Frontiers Journal Series operates on a revolutionary invention, the tiered publishing system, initially addressing specific communities of scholars, and gradually climbing up to broader public understanding, thus serving the interests of the lay society, too.

Dedication to Quality

Each Frontiers article is a landmark of the highest quality, thanks to genuinely collaborative interactions between authors and review editors, who include some of the world's best academicians. Research must be certified by peers before entering a stream of knowledge that may eventually reach the public - and shape society; therefore, Frontiers only applies the most rigorous and unbiased reviews. Frontiers revolutionizes research publishing by freely delivering the most outstanding research, evaluated with no bias from both the academic and social point of view. By applying the most advanced information technologies, Frontiers is catapulting scholarly publishing into a new generation.

What are Frontiers Research Topics?

Frontiers Research Topics are very popular trademarks of the Frontiers Journals Series: they are collections of at least ten articles, all centered on a particular subject. With their unique mix of varied contributions from Original Research to Review Articles, Frontiers Research Topics unify the most influential researchers, the latest key findings and historical advances in a hot research area! Find out more on how to host your own Frontiers Research Topic or contribute to one as an author by contacting the Frontiers Editorial Office: frontiersin.org/about/contact

RECENT INSIGHTS INTO THE DOUBLE ROLE OF HYDROGEN PEROXIDE IN PLANTS

Topic Editors:

Naser A. Anjum, Aligarh Muslim University, India

Sarvajeet Singh Gill, Maharshi Dayanand University, India

Francisco J. Corpas, Department of Biochemistry, Cell and Molecular Biology of Plants, Experimental Station of Zaidín, Spanish National Research Council (CSIC), Spain

Cristina Ortega-Villasante, Autonomous University of Madrid, Spain

Luis E. Hernandez, Autonomous University of Madrid, Spain

Narendra Tuteja, International Centre for Genetic Engineering and Biotechnology (India), India

Adriano Sofo, University of Basilicata, Italy

Mirza Hasanuzzaman, Sher-e-Bangla Agricultural University, Bangladesh

Masayuki Fujita, Kagawa University, Japan

Citation: Anjum, N. A., Gill, S. S., Corpas, F. J., Ortega-Villasante, C., Hernandez, L. E., Tuteja, N., Sofo, A., Hasanuzzaman, M., Fujita, M., eds. (2022). Recent Insights Into the Double Role of Hydrogen Peroxide in Plants. Lausanne: Frontiers Media SA. doi: 10.3389/978-2-88974-524-1

Table of Contents

- 05 Editorial: Recent Insights Into the Double Role of Hydrogen Peroxide in Plants**
Naser A. Anjum, Sarvajeet Singh Gill, Francisco J. Corpas, Cristina Ortega-Villasante, Luis E. Hernandez, Narendra Tuteja, Adriano Sofo, Mirza Hasanuzzaman and Masayuki Fujita
- 09 Hydrogen Peroxide Priming Modulates Abiotic Oxidative Stress Tolerance: Insights From Ros Detoxification And Scavenging**
Mohammad A. Hossain, Soumen Bhattacharjee, Saed-Moucheshi Armin, Pingping Qian, Wang Xin, Hong-Yu Li, David J. Burritt, Masayuki Fujita and Lam-Son P. Tran
- 28 Hydrogen Peroxide Alleviates Nickel-Inhibited Photosynthetic Responses Through Increase in Use-Efficiency of Nitrogen and Sulfur, and Glutathione Production in Mustard**
M. I. R. Khan, Nafees A. Khan, Asim Masood, Tasir S. Per and Mohd Asgher
- 48 Different Modes of Hydrogen Peroxide Action During Seed Germination**
Łukasz Wojtyła, Katarzyna Lechowska, Szymon Kubala and Małgorzata Garnczarska
- 64 Hydrogen Peroxide Signaling in Plant Development and Abiotic Responses: Crosstalk with Nitric Oxide and Calcium**
Lijuan Niu and Weibiao Liao
- 78 Identification and Comparative Analysis of H₂O₂-Scavenging Enzymes (Ascorbate Peroxidase and Glutathione Peroxidase) in Selected Plants Employing Bioinformatics Approaches**
Ibrahim I. Ozyigit, Ertugrul Filiz, Recep Vatansever, Kuaybe Y. Kurtoglu, Ibrahim Koc, Münir X. Öztürk and Naser A. Anjum
- 101 Hydrogen Peroxide, Signaling in Disguise During Metal Phytotoxicity**
Ann Cuypers, Sophie Hendrix, Rafaela Amaral dos Reis, Stefanie De Smet, Jana Deckers, Heidi Gielen, Marijke Jozefczak, Christophe Loix, Hanne Vercampt, Jaco Vangronsveld and Els Keunen
- 126 Cross Talk Between H₂O₂ and Interacting Signal Molecules Under Plant Stress Response**
Ina Saxena, Sandhya Srikanth and Zhong Chen
- 142 Brassinosteroid Ameliorates Zinc Oxide Nanoparticles-Induced Oxidative Stress by Improving Antioxidant Potential and Redox Homeostasis in Tomato Seedling**
Mengqi Li, Golam J. Ahammed, Caixia Li, Xiao Bao, Jingquan Yu, Chunlei Huang, Hanqin Yin and Jie Zhou
- 155 Tuning of Redox Regulatory Mechanisms, Reactive Oxygen Species and Redox Homeostasis Under Salinity Stress**
M. Sazzad Hossain and Karl-Josef Dietz

- 170** *A Different Pattern of Production and Scavenging of Reactive Oxygen Species in Halophytic Eutrema salsugineum (Thellungiella salsuginea) Plants in Comparison to Arabidopsis thaliana and Its Relation to Salt Stress Signaling*
 Maria Pilarska, Monika Wiciarz, Ivan Jajić, Małgorzata Kozieradzka-Kiszkurno, Petre Dobrev, Radomíra Vanková and Ewa Niewiadomska
- 182** *Patterns of ROS Accumulation in the Stigmas of Angiosperms and Visions into Their Multi-Functionality in Plant Reproduction*
 Adoración Zafra, Juan D. Rejón, Simon J. Hiscock and Juan de Dios Alché
- 189** *Involvement of Polyamine Oxidase-Produced Hydrogen Peroxide During Coleorhiza-Limited Germination of Rice Seeds*
 Bing-Xian Chen, Wen-Yan Li, Yin-Tao Gao, Zhong-Jian Chen, Wei-Na Zhang, Qin-Jian Liu and Zhuang Chen
- 202** *Corrigendum: Involvement of Polyamine Oxidase-Produced Hydrogen Peroxide During Coleorhiza-Limited Germination of Rice Seeds*
 Bing-Xian Chen, Wen-Yan Li, Yin-Tao Gao, Zhong-Jian Chen, Wei-Na Zhang, Qin-Jian Liu and Zhuang Chen
- 203** *Hydrogen Peroxide and Polyamines Act as Double Edged Swords in Plant Abiotic Stress Responses*
 Kamala Gupta, Atreyee Sengupta, Mayukh Chakraborty and Bhaskar Gupta
- 222** *Reactive Oxygen Species (ROS): Beneficial Companions of Plants' Developmental Processes*
 Rachana Singh, Samiksha Singh, Parul Parihar, Rohit K. Mishra, Durgesh K. Tripathi, Vijay P. Singh, Devendra K. Chauhan and Sheo M. Prasad
- 241** *Abscisic Acid-Induced H₂O₂ Accumulation Enhances Antioxidant Capacity in Pumpkin-Grafted Cucumber Leaves Under Ca(NO₃)₂ Stress*
 Sheng Shu, Pan Gao, Lin Li, Yinghui Yuan, Jin Sun and Shirong Guo
- 251** *ROS-Mediated Inhibition of S-nitrosoglutathione Reductase Contributes to the Activation of Anti-oxidative Mechanisms*
 Izabella Kovacs, Christian Holzmeister, Markus Wirtz, Arie Geerlof, Thomas Fröhlich, Gaby Römling, Gitto T. Kuruthukulangarakoola, Eric Linster, Rüdiger Hell, Georg J. Arnold, Jörg Durner and Christian Lindermayr
- 268** *ROS Production and Scavenging Under Anoxia and Re-Oxygenation in Arabidopsis Cells: A Balance Between Redox Signaling and Impairment*
 Annalisa Paradiso, Sofia Caretto, Antonella Leone, Anna Bove, Rossella Nisi and Laura De Gara
- 279** *Glutathionylation of Pea Chloroplast 2-Cys Prx and Mitochondrial Prx IIF Affects Their Structure and Peroxidase Activity and Sulfiredoxin Deglutathionylates Only the 2-Cys Prx*
 Aingeru Calderón, Alfonso Lázaro-Payo, Iván Iglesias-Baena, Daymi Camejo, Juan J. Lázaro, Francisca Sevilla and Ana Jiménez
- 291** *Hydrogen Peroxide Pretreatment Mitigates Cadmium-Induced Oxidative Stress in Brassica napus L.: An Intrinsic Study on Antioxidant Defense and Glyoxalase Systems*
 Mirza Hasanuzzaman, Kamrun Nahar, Sarvajeet S. Gill, Hesham F. Alharby, Bam H. N. Razafindrabe and Masayuki Fujita



Editorial: Recent Insights Into the Double Role of Hydrogen Peroxide in Plants

Naser A. Anjum^{1*†}, Sarvajeet Singh Gill^{2*†}, Francisco J. Corpas³,
Cristina Ortega-Villasante⁴, Luis E. Hernandez⁵, Narendra Tuteja⁶, Adriano Sofo⁷,
Mirza Hasanuzzaman⁸ and Masayuki Fujita⁹

¹ Department of Botany, Aligarh Muslim University, Aligarh, India, ² Stress Physiology and Molecular Biology Lab, Centre for Biotechnology, MD University, Rohtak, India, ³ Department of Biochemistry, Cell and Molecular Biology of Plants, Experimental Station of Zaidin, Spanish National Research Council (CSIC), Granada, Spain, ⁴ Fisiología Vegetal (Plant Physiology Laboratory), Dpto. Biología (Biology Department), Universidad Autónoma de Madrid, Madrid, Spain, ⁵ Laboratory of Plant Physiology, Department of Biology/Research Centre for Biodiversity and Global Change, Universidad Autónoma Madrid, Madrid, Spain, ⁶ Plant Molecular Biology Group, International Centre for Genetic Engineering and Biotechnology (ICGEB), New Delhi, India, ⁷ Department of European and Mediterranean Cultures: Architecture, Environment, and Cultural Heritage (DICEM), University of Basilicata, Potenza, Italy, ⁸ Department of Agronomy, Faculty of Agriculture, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh, ⁹ Laboratory of Plant Stress Responses, Department of Applied Biological Science, Faculty of Agriculture, Kagawa University, Takamatsu, Japan

Keywords: abiotic stresses, hydrogen peroxide, H₂O₂-metabolism, priming, plant stress tolerance

OPEN ACCESS

Edited and reviewed by:

Anna N. Stepanova,
North Carolina State University,
United States

*Correspondence:

Naser A. Anjum
dnaanjum@gmail.com
Sarvajeet Singh Gill
srgill14@yahoo.co.in

[†] These authors share first authorship

Specialty section:

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

Received: 25 December 2021

Accepted: 04 January 2022

Published: 28 January 2022

Citation:

Anjum NA, Gill SS, Corpas FJ,
Ortega-Villasante C, Hernandez LE,
Tuteja N, Sofo A, Hasanuzzaman M
and Fujita M (2022) Editorial: Recent
Insights Into the Double Role of
Hydrogen Peroxide in Plants.
Front. Plant Sci. 13:843274.
doi: 10.3389/fpls.2022.843274

Editorial on the Research Topic

Recent Insights Into the Double Role of Hydrogen Peroxide in Plants

Reactive oxygen species (ROS) of varied types can be yielded in plants at several primary sites (such as the chloroplast, mitochondria, and peroxisomes) under normal aerobic metabolism *via* processes including photosynthetic and respiratory electron transport chains. However, impaired oxidant-antioxidant balance and extreme growth conditions in plants are bound to cause increases in the cellular concentrations of radical and non-radical ROS such as superoxide anions (O^{2•-}), hydroxyl radical (OH•), singlet oxygen (¹O²), and hydrogen peroxide (H₂O₂). On the one hand, H₂O₂ has no unpaired electrons and is moderately reactive. Owing to its relative stability compared to other ROS and its capacity for diffusing through aquaporins in the membranes and over more considerable distances within the cell (Bienert et al., 2007), H₂O₂ acts as a stress signal transducer and contributes to numerous physiological functions in plants. On the other hand, H₂O₂ is a relatively long-lived molecule with a half-life of 1 ms, readily crosses biological membranes, and consequently can bring oxidative consequences far from the site of its formation (Neill et al., 2002; Sharma et al., 2012; Sehar et al., 2021). The Frontiers Research Topic “Recent Insights into the Double Role of Hydrogen Peroxide in Plants” highlighted the major mechanisms underlying the dual role of H₂O₂ in response to different abiotic stresses in plants. This Research Topic incorporated 19 publications, including 10 original research articles, 8 reviews, and one perspective article.

H₂O₂-METABOLISM AND H₂O₂-PRIMING ROLES IN ABIOTIC STRESS MANAGEMENT

As a potent signaling molecule H₂O₂ gets produced in routine in stressed or non-stressed conditions *via* dismutation of O^{2•-} radicals through superoxide dismutase (SOD) during electron transport in different compartments of the plant

cell, and is involved in the regulation of the plant growth, metabolism, and stress tolerance. It has also been noted that at higher concentrations in the cell during oxidative stress, ROS, including H₂O₂, can oxidize vital biomolecules (like nucleic acids, proteins, and lipids) and significantly impacts the seed germination process (Wojtyła et al.). Among the major abiotic stress factors, several heavy metals provoke increases in the production of ROS through plasma membrane-bound NADPH oxidases. However, the relationship of H₂O₂ has also been established in heavy metal tolerance in crop plants (Cuypers et al.). H₂O₂ directly mediates metal-induced oxidative signaling, where the production of H₂O₂ may involve H₂O₂ receptors, redox-sensitive transcription factors and inhibition of phosphatases (Miller et al., 2008). H₂O₂ sensing in metal-exposed plants also involves activation of mitogen-activated protein kinase (MAPK) pathways (Opdenakker et al., 2012). Additionally, interaction of H₂O₂ with Ca²⁺ (Baliardini et al., 2015), NO (Arasimowicz-Jelonek et al., 2012) and oxylipins (Tamás et al., 2009; Keunen et al., 2013) was also reported in metal-exposed plants. Though excess accumulation of H₂O₂ and polyamines (PAs) can be detrimental for the plant cell leading to premature cell death, a fine-tuning of these signaling molecules (H₂O₂ and PAs) can result in stress management by coordinating intra-cellular and systemic signaling systems (Gupta et al.). Polyamine oxidase (PAO)-induced production of H₂O₂ was found to be involved in the coleorhiza-limited rice seed germination (Chen et al.). ROS-specific probe DCFH2-DA enabled confocal laser scanning microscopy revealed a high level of ROS in the stigma at different developmental stages (unopened flower buds, recently opened flowers, dehiscent anthers, and flowers after fertilization) of scrutinized plants (Zafra et al.).

During evolution, plants have developed an efficient ROS-scavenging system constituting an array of enzymatic (SOD; CAT, catalase; APX, ascorbate peroxidase; GR, glutathione reductase; MDHAR, monodehydroascorbate reductase; DHAR, dehydroascorbate reductase; GPX, glutathione peroxidase; GOPX, guaiacol peroxidase, and GST, glutathione-S-transferase) and non-enzymatic (AsA, ascorbic acid; GSH, glutathione; phenolic compounds, alkaloids, non-protein amino acids, and α -tocopherols) antioxidants to get rid of excessive ROS in the cell (Singh et al.). Notably, NADH oxidase (RBOH), alternative oxidase (AOX), the plastid terminal oxidase (PTOX), and the malate valve with the malate dehydrogenase isoforms are involved in maintenance of the cellular redox homeostasis under salinity stress (Hossain and Dietz). In *Arabidopsis* cell suspension cultures, anoxia stress/shock led to significant increases in H₂O₂ (and also nitric oxide, NO); however, re-oxygenation maintained the components of ROS scavenging machinery like ascorbate-glutathione (AsA-GSH) system, α -tocopherol, and eventual cell survival as result of decreased H₂O₂ (Paradiso et al.). *Eutrema salsugineum* (halophyte) and *Arabidopsis thaliana* (glycophyte) exhibited a differential pattern of accumulation and scavenging of ROS. In particular, compared to *A. thaliana* chloroplasts, *E. salsugineum* chloroplasts showed a constitutive increase and the cell's steady-state regulation of H₂O₂ level which prepared this plant for ROS-control mainly due to an efficient ROS-scavenging machinery including glucosinolates content and

well-coordinated tuning of hormonal signaling (Pilarska et al.). Elevation in the cellular level of H₂O₂ and its consequences can be controlled by brassinosteroids, a class of plant-specific essential steroid hormones. To this end, in tomato seedlings, brassinosteroid (24-epibrassinolide) ameliorated the impacts of zinc oxide nanoparticles-caused elevated H₂O₂ by enhancing the activity of enzyme involved in superoxide-dismutation (SOD), H₂O₂-metabolizing enzymes (catalase, CAT; and APX), increasing GSH-regeneration (as a result of increased GSH reductase activity; and consequently decreasing GSH-oxidation), finally inducing the transcripts of *Cu/Zn SOD*, *GSH1*, *CAT1*, and *GRI* (Li et al.). In a comprehensive *in silico* study, APX and GSH-peroxidase (GPX) genes/proteins from 18 different plant species were identified and compared in order to unravel their significance in excessive H₂O₂ management (Ozyigit et al.). Notably, APX and GPX were found to be involved in the metabolism of antioxidants and secondary metabolites, redox homeostasis, stress adaptation, and photosynthesis/respiration. The major redox proteins namely plant peroxiredoxins (Prxs) and sulfiredoxins (Srxs) are involved in antioxidant defense and redox signaling in stressed plants. Srxs were also found to be involved in antioxidant defense and redox signaling in response to environmental stimuli; post-translational modifications of Srxs regulate the ROS-transduction and bioactivity. On the other hand, Prxs are sensitive to glutathionylation. Investigation of the glutathionylation of recombinant chloroplastic 2-Cys Prx and mitochondrial Prx IIF of pea plants revealed glutathionylation-mediated change of the decameric form of 2-Cys Prx into its dimeric glutathionylated form. Additionally, the reduced dimeric form of Prx IIF was glutathionylated without changing its oligomeric state (Calderón et al.). Thus, glutathionylation was argued to depend on the GSH/GSSG ratio owing to the perceptible difference in the exact effect on the 2-Cys Prx and Prx IIF proteins.

H₂O₂-priming (exposure of seeds, seedlings, or plants to stressors/chemical compounds that makes them ready to tolerate the later stress events) helps in biotic and abiotic stress tolerance in various crop plants by triggering the ROS scavenging machinery (Dikilitas et al., 2020). Exogenous supply of H₂O₂ can induce stress tolerance under salt, drought, chilling, high temperatures, and heavy metal stress (Hossain et al.). In a study on mustard (*Brassica juncea* L.) cultivars, H₂O₂-induced reversal of the major negative impacts of Ni stress (200 mg Ni kg⁻¹ soil) led to increased photosynthetic nitrogen-use efficiency, sulfur-use efficiency, and GSH content and decreased levels of lipid peroxidation and electrolyte leakage (Khan et al.). Notably, H₂O₂ priming-mediated increased tolerance to cadmium-caused oxidative stress in *Brassica napus* involved fine-tuning between the glyoxalase system and the components of ROS-scavenging machinery (Hasanuzzaman et al.).

H₂O₂ CROSSTALK WITH OTHER MOLECULES

Along with H₂O₂, other signaling molecules (such as nitric oxide, NO; and calcium, Ca²⁺) and phytohormones (such as jasmonic

acid, JA; salicylic acid, SA; and abscisic acid, ABA) play key roles in stress signaling cascades and crosstalk during plants' stress responses (Saxena et al., 2016). To this end, the crosstalk of H₂O₂ with NO and Ca²⁺ was argued to contribute to regulation of the plant development and abiotic stress responses (Niu and Liao). Notably, the role of SA in adventitious root formation involved H₂O₂ acting as a downstream messenger (Yang et al., 2013). Having emerged as a master regulator of stress responses, ABA signaling pathway triggers significant changes in gene expression and plants' adaptive physiological responses (Saxena et al., 2016). There occurs a close relation among the MAPK cascades, ABA, JA, SA, and H₂O₂ where exogenous application of H₂O₂ triggers MAPK cascade, which in turn involves ABA, JA, and SA (Saxena et al.). ABA-induced H₂O₂ accumulation can protect plant parts (such as pumpkin-grafted cucumber leaves) against Ca(NO₃)₂ via ABA/H₂O₂ signaling-led induction of ROS-scavenging machinery (Shu et al.). S-nitrosogluthione reductase (GSNOR) determines the level of S-nitrosothiol and thereby regulates NO-signaling in plants (Lindermayr, 2018; Jahnová et al., 2019). In *A. thaliana*, H₂O₂ *in vitro* led to inhibition of the activity of GSNOR and significantly changed NO-homeostasis, which in turn resulted in the activation of ROS-scavenging machinery in order to suppress the oxidative damage (Kovacs et al.).

CONCLUSIONS AND FUTURE PERSPECTIVE

In the current Research Topic “Recent insights into the double role of hydrogen peroxide in plants,” the contributions discussed

the versatile role of H₂O₂ as a signaling molecule that triggers the upregulation of the components of antioxidant defense machinery and imparts tolerance in crop plants against the variety of environmental cues. The crosstalk of H₂O₂ with other signaling molecules and phytohormones leads to signal transduction in response to various stresses and regulates plant growth, development, and stress tolerance. Therefore, further understanding on the coordination of H₂O₂ and other signaling molecules NO, Ca²⁺, MAPK, SA, and ABA can pave the way to achieving tolerance in crop plants to increasing stress conditions.

AUTHOR CONTRIBUTIONS

NA and SG prepared the first draft of the manuscript. FC, CO-V, LH, NT, AS, MH, and MF read and revised the manuscript. All authors listed approved the final version for publication.

ACKNOWLEDGMENTS

NA is thankful to the Department of Biotechnology (DBT), Govt. of India for the financial assistance in the form of DBT-Ramalingaswami Re-Entry Fellowship (BT/HRD/35/02/2006; BT/RLF/Re-Entry/18/2015). SG acknowledges the partial financial support received from the University Grants Commission (UGC), Council of Scientific and Industrial Research (CSIR), and Science and Engineering Research Board, Govt. of India, New Delhi.

REFERENCES

- Arasimowicz-Jelonek, M., Floryszak-Wieczorek, J., Deckert, J., Rucińska-Sobkowiak, R., Gzyl, J., Pawlak-Sprada, S., et al. (2012). Nitric oxide implication in cadmium-induced programmed cell death in roots and signaling response of yellow lupine plants. *Plant Physiol. Biochem.* 58, 124–134. doi: 10.1016/j.plaphy.2012.06.018
- Baliardini, C., Meyer, C.-L., Salis, P., Saumitou-Laprade, P., and Verbruggen, N. (2015). CATION EXCHANGER1 cosegregates with cadmium tolerance in the metal hyperaccumulator *Arabidopsis halleri* and plays a role in limiting oxidative stress in *Arabidopsis* spp. *Plant Physiol.* 169, 549–559. doi: 10.1104/pp.15.01037
- Bienert, G. P., Möller, A. L., Kristiansen, K. A., Schulz, A., Möller, I. M., Schjoerring, J. K., et al. (2007). Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J. Biol. Chem.* 282, 1183–1192. doi: 10.1074/jbc.M603761200
- Dikilitas, M., Simsek, E., and Roychoudhury, A. (2020). “Modulation of abiotic stress tolerance through hydrogen peroxide,” in *Protective Chemical Agents in the Amelioration of Plant Abiotic Stress: Biochemical and Molecular Perspectives*, eds A. Roychoudhury and D. K. Tripathi (New York, NY: John Wiley & Sons), 147–173. doi: 10.1002/9781119552154.ch7
- Jahnová, J., Luhová, L., and Petrivalský, M. (2019). S-nitrosogluthione reductase - the master regulator of protein S-nitrosation in plant NO-signaling. *Plants* 8:48. doi: 10.3390/plants8020048
- Keunen, E., Remans, T., Opendakker, K., Jozefczak, M., Gielen, H., Guisez, Y., et al. (2013). A mutant of the *Arabidopsis thaliana* LIPOXYGENASE1 gene shows altered signalling and oxidative stress related responses after cadmium exposure. *Plant Physiol. Biochem.* 63, 272–280. doi: 10.1016/j.plaphy.2012.12.005
- Lindermayr, C. (2018). Crosstalk between reactive oxygen species and nitric oxide in plants: key role of S-nitrosogluthione reductase. *Free Radic. Biol. Med.* 122, 110–115. doi: 10.1016/j.freeradbiomed.2017.11.027
- Miller, G., Shulaev, V., and Mittler, R. (2008). Reactive oxygen signaling and abiotic stress. *Physiol. Plant.* 133, 481–489. doi: 10.1111/j.1399-3054.2008.01090.x
- Neill, S., Desikan, R., and Hancock, J. (2002). Hydrogen peroxide signalling. *Curr. Opin. Plant Biol.* 5, 388–395. doi: 10.1016/S1369-5266(02)00282-0
- Opendakker, K., Remans, T., Vangronsveld, J., and Cuypers, A. (2012). Mitogen-activated protein (MAP) kinases in plant metal stress: regulation and responses in comparison to other biotic and abiotic stresses. *Int. J. Mol. Sci.* 13, 7828–7853. doi: 10.3390/ijms13067828
- Saxena, I., Srikanth, S., and Chen, Z. (2016). Cross talk between H₂O₂ and interacting signal molecules under plant stress response. *Front. Plant Sci.* 7:570. doi: 10.3389/fpls.2016.00570
- Sehar, Z., Jahan, B., Masood, A., Anjum, N. A., and Khan, N. A. (2021). Hydrogen peroxide potentiates defense system in presence of sulfur to protect chloroplast damage and photosynthesis of wheat under drought stress. *Physiol. Plant* 172, 922–934. doi: 10.1111/ppl.13225
- Sharma, P., Jha, A. B., Dubey, R. S., and Pessarakli, M. (2012). Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *J. Bot.* doi: 10.1155/2012/217037
- Tamás, L., Dudíková, J., Durceková, K., Halusková, L., Huttová, J., and Mistrík, I. (2009). Effect of cadmium and temperature on the lipoxygenase activity in barley root tip. *Protoplasma* 235, 17–25. doi: 10.1007/s00709-008-0027-2
- Yang, W., Zhu, C., Ma, X., Li, G., Gan, L., Ng, D., et al. (2013). Hydrogen peroxide is a second messenger in the salicylic acid-triggered

adventitious rooting process in mung bean seedlings. *PLoS ONE* 8:e84580. doi: 10.1371/journal.pone.0084580

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in

this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Anjum, Gill, Corpas, Ortega-Villasante, Hernandez, Tuteja, Sofo, Hasanuzzaman and Fujita. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.